

PAE Bibliography



The following list includes published journal articles regarding prostatic artery embolization (PAE).

1. Amouyal, G., Pellerin, O., Del Giudice, C., et al. (2017). Bilateral arterial embolization of the prostate through a single prostatic artery: a case series. *Cardiovasc Intervent Radiol*, May; 40(5):780-787. <https://www.ncbi.nlm.nih.gov/pubmed/27999920>
2. Amouyal, G., Chague, P., Pellerin, O., et al. (2016). Safety and efficacy of occlusion of large extra-prostatic anastomoses during prostatic artery embolization for symptomatic BPH. *Cardiovasc Intervent Radiol*, Sep;39(9):1245-1255. <https://www.ncbi.nlm.nih.gov/pubmed/27412754>
3. Amouyal, G., Thiounn, N., Pellerin, O., et al. (2016). Clinical results after prostatic artery embolization using the PErFecTED technique: a single-center study. *Cardiovasc Intervent Radiol*, Mar;39(3):367-375. <https://www.ncbi.nlm.nih.gov/pubmed/26702619>
4. Antunes, A. A., Carnevale, F. C., da Motta Leal Filho, J. M., et al. (2013). Clinical, laboratorial, and urodynamic findings of prostatic artery embolization for the treatment of urinary retention related to benign prostatic hyperplasia. A prospective single-center pilot study. *Cardiovasc Intervent Radiol*, Aug;36(4):978-986. <https://www.ncbi.nlm.nih.gov/pubmed/23580116>
5. Bhatia, S., Harward, S. H., Sinha, V. K., et al. (2017). Prostate artery embolization via transradial or transulnar versus transfemoral arterial access: technical results. *J Vasc Interv Radiol*, Jun;28(6):898-905. <https://www.ncbi.nlm.nih.gov/pubmed/28396191>
6. Bhatia, S., Sinha, V. K., Bordegaray, M., et al. (2017). Role of coil embolization during prostatic artery embolization: incidence, indications, and safety profile. *J Vasc Interv Radiol*, May;28(5):656-664. <https://www.ncbi.nlm.nih.gov/pubmed/28284886>
7. Bhatia, S., Kava, B., Pereira, K., et al. (2015). Prostate artery embolization for giant prostatic hyperplasia. *J Vasc Interv Radiol*, Oct;26(10):1583-1585. <https://www.ncbi.nlm.nih.gov/pubmed/26408223>
8. Brook, O. R., Faintuch, S., Brook, A., et al. (2013). Embolization therapy for benign prostatic hyperplasia: influence of embolization particle size on gland perfusion. *J Magn Reson Imaging*, Aug;38(2):380-387. <https://www.ncbi.nlm.nih.gov/pubmed/23239260>
9. Camara-Lopes, G., Mattedi, R., Antunes, A. A., et al. (2013). The histology of prostate tissue following prostatic artery embolization for the treatment of benign prostatic hyperplasia. *Int Braz J Urol*, Mar-Apr;39(2):222-227. <https://www.ncbi.nlm.nih.gov/pubmed/23683668>
10. Carnevale, F. C., Moreira, A. M., Harward, S. H., et al. (2017). Recurrence of lower urinary tract symptoms following prostate artery embolization for benign hyperplasia: single center experience comparing two techniques. *Cardiovasc Intervent Radiol*, Mar;40(3):366-374. <https://www.ncbi.nlm.nih.gov/pubmed/28074310>
11. Carnevale, F. C., Iscaife, A., Yoshinaga, E. M., et al. (2016). Transurethral resection of the prostate (TURP) versus original and PErFecTED prostate artery embolization (PAE) due to benign prostatic hyperplasia (BPH): preliminary results of a single center, prospective, urodynamic-controlled analysis. *Cardiovasc Intervent Radiol*, Jan;39(1):44-52. <https://www.ncbi.nlm.nih.gov/pubmed/26506952>
12. Carnevale, F. C., Moreira, A. M., & Antunes, A. A. (2014). The "PErFecTED technique": proximal embolization first, then embolize distal for benign hyperplasia. *Cardiovasc Intervent Radiol*, Dec;37(6):1602-1605. <https://www.ncbi.nlm.nih.gov/pubmed/24943914>
13. Carnevale, F. C., da Motta-Leal-Filho, J. M., Antunes, A. A., et al. (2013). Quality of life and clinical symptom improvement support prostatic artery embolization for patients with acute urinary retention caused by benign prostatic hyperplasia. *J Vasc Interv Radiol*, Apr;24(4):535-542. <https://www.ncbi.nlm.nih.gov/pubmed/23434084>
14. Carnevale, F. C. & Antunes, A. A. (2013). Prostatic artery embolization for enlarged prostates due to benign prostatic hyperplasia. How I do it. *Cardiovasc Intervent Radiol*, Dec;36(6):1452-1463. <https://www.ncbi.nlm.nih.gov/pubmed/23903785>
15. Carnevale, F. C., da Motta-Leal-Filho, J. M., Antunes, A. A., et al. (2011). Midterm follow-up after prostate embolization in two patients with benign prostatic hyperplasia. *Cardiovasc Intervent Radiol*, Dec;34(6):1330-1333. <https://www.ncbi.nlm.nih.gov/pubmed/21387120>
16. Carnevale, F. C., Antunes, A. A., da Motta Leal Filho, J. M., et al. (2010). Prostatic artery embolization as a primary treatment for benign prostatic hyperplasia: preliminary results in two patients. *Cardiovasc Intervent Radiol*, Apr;33(2):355-361. <https://www.ncbi.nlm.nih.gov/pubmed/19908092>
17. de Assis, A. M., Moreira, A. M., de Paula Rodrigues, V. C., et al. (2015). Pelvic arterial anatomy relevant to prostatic artery embolization and proposal for angiographic classification. *Cardiovasc Intervent Radiol*, Aug;38(4):855-861. <https://www.ncbi.nlm.nih.gov/pubmed/25962991>
18. de Assis, A. M., Moreira, A. M., de Paula Rodrigues, V. C., et al. (2015). Prostatic artery embolization for treatment of benign prostatic hyperplasia in patients with prostates > 90 g: a prospective single-center study. *J Vasc Interv Radiol*, Jan;26(1):87-93. <https://www.ncbi.nlm.nih.gov/pubmed/25541446>
19. Feng, S., Tian, Y., Liu, W., et al. (2017). Prostatic arterial embolization treating moderate-to-severe lower urinary tract symptoms related to benign prostatic hyperplasia: a meta-analysis. *Cardiovasc Intervent Radiol*, Jan;40(1):22-32. <https://www.ncbi.nlm.nih.gov/pubmed/27872988>
20. Frenk, N. E., Baroni, R. H., Carnevale, F. C., et al. (2014). MRI findings after prostatic artery embolization for treatment of benign hyperplasia. *AJR Am J Roentgenol*, Oct;203(4):813-821. <https://www.ncbi.nlm.nih.gov/pubmed/25247946>
21. Gabr, A. H., Gabr, M. F., Elmohamady, B. N., et al. (2016). Prostatic artery embolization: a promising technique in the treatment of high-risk patients with benign prostatic hyperplasia. *Urol Int*, Oct;97(3):320-324. <https://www.ncbi.nlm.nih.gov/pubmed/27322582>
22. Golzarian, J., Antunes, A. A., Bilhim, T., et al. (2014). Prostatic artery embolization to treat lower urinary tract symptoms related to benign prostatic hyperplasia and bleeding in patients with prostate cancer: proceedings from a multidisciplinary research consensus panel. *J Vasc Interv Radiol*, May;25(5):665-674. <https://www.ncbi.nlm.nih.gov/pubmed/24560898>
23. Gonçalves, O. M., Carnevale, F. C., Moreira, A. M., et al. Comparative study using 100-300 versus 300-500 µm microspheres for symptomatic patients due to enlarged-BPH prostates. *Cardiovasc Intervent Radiol*, Oct;39(10):1372-1378. <https://www.ncbi.nlm.nih.gov/pubmed/27509889>
24. Grosso, M., Balderi, A., Arnò, M., et al. (2015). Prostatic artery embolization in benign prostatic hyperplasia: preliminary results in 13 patients. *Radiol Med*, Apr;120(4):361-368. <https://www.ncbi.nlm.nih.gov/pubmed/25245496>
25. Isaacson, A. J., Fischman, A. M., & Burke, C. T. (2016). Technical feasibility of prostatic artery embolization from a transradial approach. *AJR Am J Roentgenol*, Feb;206(2):442-444. <https://www.ncbi.nlm.nih.gov/pubmed/26797376>
26. Isaacson, A. J., Raynor, M. C., Yu, H., et al. (2016). Prostatic artery embolization using Embosphere Microspheres for prostates measuring 80-150 cm³: early results from a US trial. *J Vasc Interv Radiol*, May;27(5):709-714. <https://www.ncbi.nlm.nih.gov/pubmed/27106645>
27. Kably, I., Pereira, K., Chong, W., et al. (2016). Prostate artery embolization (PAE) in the management of refractory hematuria of prostatic origin secondary to iatrogenic urological trauma: a safe and effective technique. *Urology*, Feb;88:218-221. <https://www.ncbi.nlm.nih.gov/pubmed/26610676>
28. Kurbatov, D., Russo, G. I., Lepetukhin, A., et al. (2014). Prostatic artery embolization for prostate volume greater than 80 cm³: results from a single-center prospective study. *Urology*, Aug;84(2):400-404. <https://www.ncbi.nlm.nih.gov/pubmed/24929948>
29. Laborda, A., de Assis, A. M., Ioakeim, I., et al. (2015). Radiodermatitis after prostatic artery embolization: case report and review of the literature. *Cardiovasc Intervent Radiol*, Jun;38(3):755-759. <https://www.ncbi.nlm.nih.gov/pubmed/25799951>
30. Lin, Y. T., Amouyal, G., Thiounn, N., et al. Intra-vesical prostatic protrusion (IPP) can be reduced by prostatic artery embolization. *Cardiovasc Intervent Radiol*, May;39(5):690-695. <https://www.ncbi.nlm.nih.gov/pubmed/26542027>
31. Leite, L. C., de Assis, A. M., Moreira, A. M., et al. (2017). Prostatic tissue elimination after prostatic artery embolization (PAE): a report of three cases. *Cardiovasc Intervent Radiol*, Jun;40(6):937-941. <https://www.ncbi.nlm.nih.gov/pubmed/28175975>
32. McWilliams, J. P., Kuo, M. D., Rose, S. C., et al. (2014). Society of Interventional Radiology position statement: prostate artery embolization for treatment of benign disease of the prostate. *J Vasc Interv Radiol*, Sep;25(9):1349-1351. <https://www.ncbi.nlm.nih.gov/pubmed/24993818>

33. Moreira, A. M., Marques, C. F., Antunes, A. A., et al. (2013). Transient ischemic rectitis as a potential complication after prostatic artery embolization: case report and review of the literature. *Cardiovasc Intervent Radiol*, Dec;36(6):1690-1694. <https://www.ncbi.nlm.nih.gov/pubmed/24091752>

34. Rampoldi, A., Barbosa, F., Secco, S., et al. (2017). Prostatic artery embolization as an alternative to indwelling bladder catheterization to manage benign prostatic hyperplasia in poor surgical candidates. *Cardiovasc Intervent Radiol*, Apr;40(4):530-536. <https://www.ncbi.nlm.nih.gov/pubmed/28130568>

35. Russo, G. I., Kurbatov, D., Sansalone, S., et al. (2015). Prostatic arterial embolization vs open prostatectomy: a 1-year matched-pair analysis of functional outcomes and morbidities. *Urology*, Aug;86(2):343-348. <https://www.ncbi.nlm.nih.gov/pubmed/26199151>

36. Sun, F., Crisóstomo, V., Báez-Díaz, C., et al. (2016). Prostatic artery embolization (PAE) for symptomatic benign prostatic hyperplasia (BPH): part 2, insights into the technical rationale. *Cardiovasc Intervent Radiol*, Feb;39(2):161-169. <https://www.ncbi.nlm.nih.gov/pubmed/26563245>

37. Sun, F., Crisóstomo, V., Báez-Díaz, C., et al. (2016). Prostatic artery embolization (PAE) for symptomatic benign prostatic hyperplasia (BPH): part 1, pathological background and clinical implications. *Cardiovasc Intervent Radiol*, Jan;39(1):1-7. <https://www.ncbi.nlm.nih.gov/pubmed/26581418>

38. Sun, F., Sánchez, F. M., Crisóstomo, V., et al. Transarterial prostatic embolization: initial experience in a canine model. *AJR Am J Roentgenol*, Aug;197(2):495-501. <https://www.ncbi.nlm.nih.gov/pubmed/21785100>

39. Teoh, J. Y., Chiu, P. K., Yee, C. H., et al. (2017). Prostatic artery embolization in treating benign prostatic hyperplasia: a systematic review. *Int Urol Nephrol*, Feb;49(2):197-203. <https://www.ncbi.nlm.nih.gov/pubmed/27896579>

40. Uflacker, A., Haskal, Z. J., Bilhim, T., et al. (2016). Meta-analysis of prostatic artery embolization for benign prostatic hyperplasia. *J Vasc Interv Radiol*, Nov;27(11):1686-1697. <https://www.ncbi.nlm.nih.gov/pubmed/27742235>

Embosphere PRO™

Prostatic Artery Embolization Kit

PAE is a technically challenging procedure that demands a predictable, targeted and established microsphere like Embosphere® Microspheres. Merit Medical's new Embosphere PRO™ Prostatic Artery Embolization Kit combines the power and predictability of Embosphere® Microspheres with basic preparation and delivery tools for added convenience.

Each kit includes:

- (1) prefilled syringe of Embosphere® Microspheres, 300-500 µm or 100-300 µm
- (1) 10 mL Medallion® Syringe
- (1) 3 mL Medallion® Syringe
- (2) 1 mL Medallion® Syringes
- (1) 3-way Marquis® Stopcock, 1050 psi



This bibliography is a partial list of published journal articles; it is not all inclusive.
*Paid consultants of Merit Medical.

Before using refer to Instructions for Use for indications, contraindications, warnings, precautions, and directions for use.



Understand. Innovate. Deliver.™

Merit Medical Systems, Inc.
1600 West Merit Parkway
South Jordan, Utah 84095
1.801.253.1600
1.800.35.MERIT

Merit Medical Europe, Middle East, & Africa (EMEA)
Amerikalaan 42, 6199 AE Maastricht-Airport
The Netherlands
+31 43 358 82 22

Merit Medical Ireland Ltd.
Parkmore Business Park West
Galway, Ireland
+353 (0) 91 703 733

merit.com/pae

Australia
+61 (0) 3 8373 4943

Austria
0800 295 374

Belgium
0800 72 906 (Dutch)
0800 73 172 (Français)

Brazil
+55 11 4561 - 0062

Canada
1.800.364.4370

China
+86-10 8561 0788

Denmark
80 88 00 24

Finland
0800 770 586

France
0800 91 60 30

Germany
0800 182 0871

Hong Kong
+852-2207 6888

India
+91-80-41223376

Ireland (Republic)
1800 553 163

Italy
800 897 005

Japan
800 897 005

Luxembourg
8002 25 22

Netherlands
0800 022 81 84

Norway
800 11629

Portugal
308 801 034

Russia
+7 495 221 89 02

Singapore
+65 67536311

South Korea
+8210 2205 0707

Spain
+34 911238406

Sweden
020 792 445

Switzerland
(Deutsch)
+41 22 518 02 30

(Français)
+41 22 518 02 52

(Italiano)
+41 22 518 00 35

UK
0800 973 115